

the health implications of climate change and the science that will help us adapt to these challenges.

The authors declare they have no competing financial interests.

Roger Glass
Joshua Rosenthal
Christine M. Jessup

Fogarty International Center
National Institutes of Health
Department of Health and Human Services
Bethesda, Maryland
E-mail: joshua_rosenthal@nih.gov

Linda Birnbaum
Chris Portier

National Institute of Environmental
Health Sciences
National Institutes of Health
Department of Health and Human Services
Research Triangle Park, North Carolina

REFERENCES

- ARRA (American Recovery and Reinvestment Act). 2009. Public Law 111-5.
Ebi KL, Balbus J, Kinney PL, Lipp E, Mills D, O'Neill MS, et al. 2009. U.S. funding is insufficient to address the human health impacts and public health responses to climate variability and change. *Environ Health Perspect* 117:857–862.
NIH (National Institutes of Health). 2009a. Estimates of Funding for Various Research, Condition, and Disease Categories (RCDC). Available: <http://report.nih.gov/rcdc/categories/> [accessed 20 June 2009].
NIH (National Institutes of Health). 2009b. Recovery Act Limited Competition: NIH Challenge Grants in Health and Science Research (RC1) RFA-OD-09-003. Available: <http://grants.nih.gov/grants/guide/rfa-files/RFA-OD-09-003.html> [accessed 20 June 2009].

Editor's note: In accordance with journal policy, Ebi et al. were asked whether they wanted to respond to this letter, but they chose not to do so.

Lead Exposures from Car Batteries—A Global Problem

doi:10.1289/ehp.0901163

In “Mass Lead Intoxication from Informal Used Lead Acid Battery Recycling in Dakar, Senegal,” Haefliger et al. (2009) described a problem throughout the developing world that is both tragic and only now beginning to be understood with respect to its extent and effect.

Eighteen children (and more since) died from acute lead poisoning in late 2008 in Dakar. These poisonings occurred because the individuals recycling car batteries melted slag without appropriate controls and without having any understanding of the toxicity of lead. Most of these recyclers were women who brought their children to their work sites without knowing the risks.

These problems are not restricted to Senegal. Without much effort, investigators

from Blacksmith Institute have identified another 22 sites worldwide that are similar to this one. The identified sites are in cities in poor countries, especially in the tropics (e.g., the Dominican Republic, Philippines, Panama, El Salvador, Guatemala, India, Ghana, Jamaica) (Blacksmith Institute 2009).

Epidemiologic studies of exposed populations, such as the one in Senegal reported by Haefliger et al. (2009), are urgently needed to characterize exposures and identify related health effects. An earlier example of such a study was conducted in the Dominican Republic at Haina (also known as Bajos de Haina), which has been called the “Dominican Chernobyl.” This community is near an abandoned lead-acid battery recycling smelter, and most of the residents showed signs of lead poisoning.

The Haina site, as well as the surrounding area, was the scene of severe lead poisoning in the 1990s. In March 1997, 116 children were surveyed, and 146 children were surveyed in August 1997. Mean blood lead concentrations were 71 µg/dL (range, 9–234 µg/dL) in March and 32 µg/dL (range, 6–130 µg/dL) in August (Kaul et al. 1999). The study revealed that at least 28% of the children required immediate treatment and 5% had lead levels > 79 µg/dL, putting them at risk for severe neurologic sequelae at the time of the study. In the United States, the action level for blood lead concentration is 10 µg/dL (Centers for Disease Control and Prevention 2007; U.S. Environmental Protection Agency 2000).

The scientific findings from Haina (Kaul et al. 1999) drove a collaborative cleanup of this site, which has recently been completed. The Blacksmith Institute helped locate funding, worked closely with local authorities, and provided technical assistance to assure the cleanup was adequate. We are currently beginning a similar cleanup project in Dakar, at the site studied by Haefliger et al. (2009).

Almost all large urban centers in the developing world have a problem with recycling used lead acid batteries, and hundreds of thousands, if not millions, of children are exposed to lead from battery recycling. In humid conditions, car batteries need to be replaced every 2 or 3 years, and car use is increasing throughout the world, which will result in even more used batteries. Thus, this problem deserves our immediate and serious attention.

Blacksmith Institute, a registered 501(c)3 non-profit organization, is committed to solving pollution problems around the world. R.F. is the founder and president of Blacksmith Institute.

Richard Fuller

Blacksmith Institute
New York, New York
E-mail: fuller@blacksmithinstitute.org

REFERENCES

- Blacksmith Institute. 2009. Our Programs and Projects. Available: <http://www.blacksmithinstitute.org/our-programs-and-projects.html> [accessed 4 November 2009].
Centers for Disease Control and Prevention. 2007. Interpreting and managing blood lead levels <10 µg/dL in children and reducing childhood exposures to lead: recommendations of CDC's Advisory Committee on Childhood Lead Poisoning Prevention. *MMWR Recomm Rep* 56:1–16. Available: <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5608a1.htm> [accessed 4 November 2009].
Haefliger P, Mathieu-Nolf M, Locicero S, Ndiaye C, Coly M, Diouf A, et al. 2009. Mass lead intoxication from informal used lead acid battery recycling in Dakar, Senegal. *Environ Health Perspect* 117:1535–1540.
Kaul B, Sandhu RS, Depratt C, Reyes F. 1999. Follow-up screening of lead-poisoned children near an auto battery recycling plant, Haina, Dominican Republic. *Environ Health Perspect* 107:917–920.
U. S. Environmental Protection Agency. 2000. Concentrations of Lead in Blood. Available: http://yosemite.epa.gov/OCHP/OCHPWEB.nsf/content/blood_lead_levels.htm, [accessed 4 November 2009].

Editor's note: In accordance with journal policy, Haefliger et al. were asked whether they wanted to respond to this letter, but they chose not to do so.

Periodontal Disease and Environmental Cadmium Exposure

doi:10.1289/ehp.0901189

We were pleased to see the article by Arora et al. (2009), which describes an association between environmental exposure to cadmium and periodontal disease.

In their cross-sectional study among U.S. adults, Arora et al. (2009) found periodontal disease in 15.4% of a nationally representative sample of 11,412 participants. The authors reported that for individuals with periodontal disease, as defined in their study, the geometric mean concentration of urinary Cd (0.50 µg/g creatinine) was significantly higher than for persons with no evidence of periodontitis (0.30 µg/g creatinine).

Arora et al. (2009) correctly stated that the main source of human exposure to environmental Cd is smoking. They proposed that additional sources of Cd in the general population are “emissions from industrial activities, including mining, smelting, and manufacturing of batteries, pigments, stabilizers, and alloys” (Arora et al. 2009).

However, in our view, one Cd source has been overlooked: intraoral dental alloys. Individuals with dental alloy restorations are regularly exposed to a number of trace elements that are continuously released from intraoral alloys (Wataha 2000).

Cadmium may be released from intraoral alloys in dental patients and may be accumulated in both teeth and oral tissues, binding tightly to metallothioneins (Goyer and Clarkson 2001; Munksgaard 1992).